

Isotopic measurements of greenhouse gases at mountain sites: *Opportunities and challenges*


Eliza Harris

Climate and Environmental Physics, University of Bern *and*

Foundation for the High Altitude Research Stations Jungfrauoch and Gornergrat



Jungfraujoch and Gornergrat




- Support high-altitude scientific research, including environmental monitoring and long-term datasets



Measuring clean air


- Background and baseline stations allow us to understand large-scale changes in atmospheric processes and biogeochemical cycles



- Measurements and interest began in the 1970s following WMO's BAPMoN program, eventually becoming GAW

Mountain sites: Diverse laboratories


Jungfrauoch



Images: CSIRO (Cape Grim), AMOF (Cape Verde), NOAA (South Pole), <https://www.railbookers.com/destinations/europe/switzerland/bern/> (Thun)

Nitrogen cycle: Major research questions

- Why are N₂O emissions accelerating?
 - Increasing N inputs or climate feedbacks?
- Nitrogen losses: Changing in a changing climate?





- <2000: ~0.6 ppb / year
- >2015: ~1.0 ppb / year

Nitrogen cycle: Major research questions


- Why are N₂O emissions accelerating?
 - Increasing N inputs or climate feedbacks?
- Nitrogen losses: Changing in a changing climate?

*Global average
emission factors*




Using isotopes for biogeochemistry

- Isotopes give us information about sources, sinks and processes





Many views on the same question

- Using isotopes and other novel tracers allows us to investigate the same question from different angles:



^{14}C : Quantifying fossil fuel CO_2


^{13}C : Estimating contributions from different sources and sinks




O_2 : Quantifying fossil fuel CO_2 and oceanic and biogenic sources and sinks

Isotopes at background stations

- Isotopic timeseries are primarily from flask sample programs
- **CO₂**: Solid understanding of isotopic trends at many sites
- **CH₄**: Clear trends at some sites, particularly for $\delta^{13}\text{C}-\text{CH}_4$
- **N₂O**: Few sites, clear trend in $\delta^{15}\text{N}^{\text{bulk}}$, no spatial patterns resolved



How do we measure isotopic composition?



Images: Thermo Fisher MAT253, 2020





- **IRMS:** Separates isotopes based on mass to charge ratio – requires large, heavy instrumentation

- ✓ Precise, well understood, small sample sizes
- ✗ Not (easily) field deployable, cannot distinguish isotopomers

How do we measure isotopic composition?

- **Spectroscopy** techniques use specific absorption bands in the mid-IR to quantify different isotopocules
- ✓ Field deployable, online, quantification of isotopomers
- ✗ Interferences, corrections, drift, sample size


From Harris et al. (2014) Analytical Chemistry, 10.1021/ac403606u




Measuring isotopes at mountain sites

- Small signals, difficult logistics → calibration is challenging

Calibration and measurements simpler,
better intercomparability






Logistics simpler, measurements difficult,
better time resolution




Interpreting isotopic data from mountain sites

- Data analysis remains challenging...
 - Modelling mountain sites is always challenging
 - Source end members are poorly constrained
 - Processes are even more difficult to model
- Handling of error and uncertainty is key



Case study: N₂O isotopes at Jungfrauoch

- N₂O mixing ratio has been measured at Jungfrauoch since 2004




- **Increasing** at a rate of ~0.9 ppb / y mainly due to agriculture
- Can we see a change in the isotopic composition?
- What does this tell us about sources and sinks?

Case study: N₂O isotopes at Jungfraujoch


- Trace gases have low concentrations, eg. N₂O = 335 parts per billion in air
- *Preconcentration* is used to trap N₂O and reach >10 ppm in a laser cell



Ambient air (flask):
~335 ppb N₂O




HayeSepD:
Warmed to 45°C





Case study: N₂O isotopes at Jungfrauoch

- $\delta^{15}\text{N}$ trend reflects the light anthropogenic source





- Seasonal N₂O cycle suggested stratospheric intrusions in summer
- Isotopes agree somewhat...



Case study: N₂O isotopes at Jungfrauoch

- $\delta^{15}\text{N}$ trend reflects the light anthropogenic source



- $\delta^{15}\text{N}$ of soil can be combined with isotopic trends to map global emissions
- Increasing emissions from subtropical regions could be driving global increases

Conclusions

- Background measurements allow us to understand large-scale variability in transport, biogeochemical cycles, and anthropogenic emissions of **trace gases**
- Mixing ratios can be used to quantify total sources, but isotopes and novel tracers give us information and **sources, sinks and processes**
- Measurements can be made *in situ* or using **flask sampling**, which improves intercomparability or time resolution respectively
 - With both methods, small signals make calibration a major challenge
- Currently, isotopic timeseries are only adequate for CO₂
- *Isotopes at background sites offer many opportunities, despite many challenges!*

